


Lauric Acid

Basic Information



Lauric Acid Structure

IUPAC Name	Dodecanoic acid
CAS Number	143-07-7
HS Code	2915.90
Molecular Formula	C12H24O2
Structural Formula	CH3(CH2)10COOH
Synonyms	Dodecanoic acid, n-Dodecanoic acid, n-Lauric acid, C12 fatty acid
Molecular Weight	200.32 g/mol

Description

Lauric acid is a saturated medium-chain fatty acid with a 12-carbon chain. It is the most abundant fatty acid in coconut oil and palm kernel oil, constituting approximately 45–55% of their fatty acid composition.

Lauric acid is prized for its exceptional antimicrobial activity against bacteria, viruses, and fungi. This property drives its use in personal care products, pharmaceuticals, and food preservation.

Industrially, lauric acid is a key feedstock for producing surfactants (sodium lauryl sulfate, lauryl alcohol), emulsifiers, soaps, and metallic soaps used as PVC stabilizers. Its derivatives are widely used in detergents, cosmetics, and lubricants.

Chemical and Physical Properties

Physical Description	White crystalline powder or flakes with a characteristic soapy odor.
Color / Form	White.
Odor	Characteristic, mild soapy odor.
Taste	Bland to slightly soapy.
Boiling Point	298–299 °C
Melting Point	43–44 °C
Flash Point	~180 °C
Solubility	Practically insoluble in water; soluble in ethanol, diethyl ether, and benzene.
Density	Approx. 0.883 g/cm³ (liquid)
Vapor Density	Greater than air.
Vapor Pressure	Very low at ambient temperature.
Stability / Shelf Life	Stable under normal conditions. Resistant to oxidation.
Viscosity	Low viscosity liquid when molten (above 44 °C).
Heat of Combustion	Approx. 77,390 kJ/mol.
Polymerization	No hazardous polymerization known.

Ionization Potential **No data available.**

Uses and Manufacturing

Uses

Lauric acid is one of the most industrially important fatty acids derived from coconut and palm kernel oil. Its primary industrial use is as a feedstock for the production of sodium lauryl sulfate (SLS) and related anionic surfactants, which are among the most widely used detergent and foaming agents in shampoos, toothpastes, and cleaning products globally. As a soap-making raw material, lauric acid produces soaps with excellent lathering and cleansing characteristics — the high lather quality of coconut oil-based soaps is primarily due to their high lauric acid content. Metallic laurates (zinc laurate, calcium laurate) are used as PVC stabilizers, mold release agents in plastics manufacturing, and lubricant additives. In the personal care and cosmetics industry, lauric acid contributes antimicrobial, emollient, and cleansing properties. Monolaurin (glycerol monolaurate, derived from lauric acid) is a potent antimicrobial agent with activity against enveloped viruses, bacteria, and yeasts, and is used in food preservation, personal care preservative systems, and as a potential nutraceutical.

Food-grade lauric acid is used as a flavoring agent (it contributes a soapy, waxy note), and as a component of confectionery fats and cocoa butter substitutes. Lauric acid-rich fats are important in confectionery because lauric acid contributes a desirable sharp melting point, quick-melting texture, and resistance to bloom.

Pharmaceutical applications include its use as a penetration enhancer in transdermal drug delivery and as a lipid excipient in lipid-based drug delivery formulations.

Methods of Manufacturing

Lauric acid is produced industrially by hydrolysis (fat splitting) of coconut oil or palm kernel oil, which contain 45–55% lauric acid in their triglyceride fraction. The fat splitting is typically conducted using the Colgate-Emery continuous high-pressure steam splitting process at temperatures of 250–260 °C and pressures of 50–60 bar, or by enzymatic hydrolysis using lipase enzymes.

The crude fatty acid mixture from splitting contains C8, C10, C12, C14, C16, C18, and C18:1 fatty acids in proportions reflecting the original oil composition. Fractional distillation under vacuum is used to separate individual fatty acids, with the C12 (lauric acid) cut being isolated as a high-purity fraction.

High-purity lauric acid (>99% GC purity) is obtained by additional recrystallization from solvents or melt crystallization. Quality is characterized by GC purity, acid value, iodine value, melting point, color (APHA), and moisture content.

Hazard Identification

Hazard Summary

Low toxicity. May cause mild skin and eye irritation. Dust may irritate respiratory tract.

Fire Hazard

Combustible solid. Dust may form ignitable mixtures.

Skin, Eye & Respiratory Irritations

Mild irritation to skin and eyes on direct contact.

Safety and First Aid

Physical Dangers

Combustible solid; dust may form ignitable mixtures with air.

Skin First Aid

Wash with soap and water.

Eye First Aid

Rinse with water for 15 minutes.

Ingestion First Aid

Generally low toxicity. Seek medical advice if large amounts are ingested.

Fire Fighting Procedures

Use CO₂, foam, or dry chemical. Water may be used to cool containers.

Handling and Storage**Nonfire Spill Response**

Small spill (solid): Sweep up carefully to avoid dust generation. Collect in labeled waste containers. For molten spill, allow to solidify then collect mechanically.

Large spill: Prevent entry into drains and waterways. Bund area. Collect mechanically after solidification or pump if liquid.

Dispose according to local environmental regulations.

Safe Storage

Store in original, tightly closed containers in a cool, dry, well-ventilated area. Keep away from strong oxidizing agents, strong bases, and reactive metals. Protect from moisture and heat. Bulk melt storage in heated stainless steel or mild steel tanks at 50–60 °C. Use nitrogen blanket in bulk storage to prevent oxidation.

Storage Conditions

Recommended storage temperature: 15–30 °C (solid); 50–55 °C if stored as melt. Shelf life: 24 months in original sealed containers. Suitable containers: HDPE drums, multi-wall paper bags, stainless steel 304/316 tanks. Avoid copper, bronze, and galvanized iron containers. Protect from moisture. Conduct periodic quality checks (acid value, color).